

EFFECTS OF SODIUM SELENITE SUPPLEMENTATION ON PRE β -HIGH-DENSITY LIPOPROTEIN FORMATION-RELATED PROTEINS IN HUMAN PRIMARY HEPATOCYTES

Mirasari Putri^{1,2*}, Chiho Yamazaki¹, Mas Rizky A. A. Syamsunarno^{2,3}, Irma Melyani Puspitasari^{1,4}, Rizky Abdulah⁴, Satomi Kameo¹, Tatsuya Iso², Masahiko Kurabayashi² and Hiroshi Koyama¹

¹Department of Public Health, Gunma University Graduate School of Medicine, Gunma, Japan, ²Department of Medicine and Biological Science, Gunma University Graduate School of Medicine, Gunma, Japan, ³Department of Biochemistry, Faculty of Medicine, Universitas Padjadjaran, Jatinangor, Indonesia, ⁴Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia

*Corresponding Author: putrimirasari@yahoo.com

Received on: 26th October, 2014

Accepted on: 6th December, 2014

ABSTRACT

The effects of selenium on pre β -HDL formation were observed in a human primary hepatocyte (Hc cells) under a basal state condition, to represent the human liver in healthy conditions. The Hc cells were cultured in a medium supplemented with 0–10 μ M sodium selenite. The effects of sodium selenite supplementation on several target proteins and genes related to pre β -HDL formation were measured by western blot analysis and real-time PCR, respectively. Protein expressions of GPx-1 and apolipoprotein A-I (apoA-I) were upregulated after treatment with sodium selenite. These results were confirmed by increased mRNA expressions of these two genes. The optimum effects of sodium selenite supplementation on protein and mRNA expressions of apoA-I and GPx-1 occurred at 50 nM; however, higher concentrations reduced the effect. In contrast, the expression levels of other pre β -HDL formation-related proteins and mRNA, apolipoprotein A-II (apoA-II) and ATP-binding cassette transporter-1 (ABCA-1), were not significantly affected. These results suggest that sodium selenite supplementation might play a role in pre β -HDL formation in Hc cells under basal state at low doses but not at high doses. Thus, an appropriate dose of selenium supplementation is essential for achieving the therapeutic potential of selenium supplementation for preventing CVD events in healthy individuals.

Keywords: Cardiovascular disease, Glutathione peroxidase, High-density lipoprotein, Selenium

INTRODUCTION

Selenium is an essential trace element to human health, and it is naturally found in grains, cereal, and meat. Selenium deficiency is related to several disorders, including cardiovascular disease (CVD) (Brenneisen *et al.*, 2005). We have previously demonstrated that selenium supplementation (organic forms, inorganic forms, and selenium-enriched foods) can induce apoptosis in cancer cells, suggesting that supplementation is beneficial for prevention and treatment of cancer (Abdulah *et al.*, 2005, 2009). Glutathione peroxidase-1 (GPx-1), a selenium-dependent enzyme, acts as a major intracellular antioxidant by reducing hydrogen peroxide and lipid peroxides; further, it also acts as a peroxynitrite reductase (Blankenberg *et al.*, 2003). A deficiency in GPx-1 activity is associated with the development of cardiovascular events and adverse prognoses in patients with CVD

(Blankenberg *et al.*, 2003).

The incidence of CVD remains one of the highest among non-communicable diseases, and CVD is the highest cause of mortality worldwide (WHO, 2011). In a recent report, a low serum concentration of high-density lipoprotein (HDL) was found to be a significant and independent risk factor for CVD (van der Steeg *et al.*, 2008). Apolipoprotein A-I (apoA-I) and apolipoprotein A-II (apoA-II) are the major proteins comprising HDL; therefore, apoA levels represent HDL cholesterol levels (Ferrier and Denise, 2011; Kontush and Chapman, 2012). ApoA-I is mainly produced in the liver and small intestine and is then secreted into the blood stream. After acquiring phospholipids and free cholesterol mediated by hepatic and intestinal ATP-binding cassette transporter A1 (ABCA-1), it then becomes nascent pre β -HDL (Ferrier and Denise, 2011; Kontush and Chapman, 2012). Additional phospholipids and free cholesterol from

extrahepatic tissue are then necessary for the transformation to mature HDL particles (Kontush and Chapman, 2012).

The selenium concentration in the blood is positively correlated with a lower atherogenic index, including higher HDL concentration or apolipoprotein levels (Koyama *et al.*, 1995; Miyazaki *et al.*, 2002; Laclaustra *et al.*, 2010; Stranges *et al.*, 2011). An *in vitro* study demonstrated that selenium supplementation increased apoA-I expression in hepatoma cells cultured under lipoprotein saccharide-induced oxidative stress (Stahle *et al.*, 2009). Selenium exists in various chemical forms, and the physiological effects of the different chemical forms vary considerably. Sodium selenite, an inorganic selenium form, is used as a supplement; recently, it is being increasingly consumed by healthy people to delay the onset or progression of age-related degenerative diseases, including CVD (Actis-Goretta *et al.*, 2004). Colpo and colleagues (Colpo *et al.*, 2013) found an increase in the HDL concentration in healthy persons after consuming Brazilian nuts, one of the richest food sources of selenium. In an experimental study, the HDL concentration was increased in male Sprague-Dawley rats fed with a high fat diet containing sodium selenite (Kaur and Bansal, 2009). In contrast, another study showed no beneficial effects of selenium supplementation against CVD (Rees *et al.*, 2013). Therefore, the efficacy of sodium selenite supplementation for preventing CVD remains inconclusive.

Selenium may increase HDL production by inducing the formation of pre β -HDL-related proteins, such as apoA-I, apoA-II, and ABCA-1, through GPx-1 activity. Nuclear factor κ B (NF- κ B) is a major transcription factor that controls the expression of various genes, and it is primarily involved in immune, inflammatory, and stress responses (Panicker *et al.*, 2010). When the NF- κ B signaling cascade is suppressed, apoA and ABCA-1 expression is facilitated (Wang *et al.*, 2006; Stahle *et al.*, 2009). GPx-1 can induce the formation of pre β -HDL-related proteins through several possible mechanisms. First, GPx-1 delays I- κ B (inhibitor κ B) degradation by inhibiting the phosphorylation of I- κ B (Kretz-Remy and Arrigo, 2001), thereby suppressing the NF- κ B cascade. Second, GPx-1 inhibits the production of reactive oxygen species, which activate NF- κ B transcription (Makropoulos *et al.*, 1996). Third, GPx-1 directly inhibits the binding of p65 and the transcription factor PPAR α , leading to increased availability of PPAR α for the transcription of pre- β HDL-related genes (Stahle *et al.*, 2009).

In the present study, we observed the effects of selenium supplementation on the expression of pre β -HDL-related proteins and mRNA in human primary hepatocytes (Hc cells) under basal state conditions. We used non-malignant cells for a better representation of the human liver in healthy conditions, to mimic healthy individuals receiving selenium supplementation in an *in vitro* system. We hypothesized that selenium supplementation would induce GPx-1 expression, which would upregulate apoA and ABCA-1 expressions,

suggesting the induction of pre β -HDL formation in the normal human liver.

MATERIALS AND METHODS

CELL CULTURE

The Hc cells were kindly provided by Prof. Takeaki Nagamine, who purchased them from the Applied Cell Biology Research Institute (Kirkland, WA). The cells were maintained as described in a previous study (Hayakawa and Nagamine, 2014) and incubated on dishes coated with type I collagen. For the experiments, cells were seeded at a concentration of 1.5×10^6 cells per 100-mm dish and incubated for 0–72 h. Hc cells were cultured under the same conditions for the selenium-supplemented groups, except that sodium selenite was added to the medium at the specified concentrations.

Before each treatment, we measured the selenium concentration in FBS and determined it to be 257.65 nM. Therefore, the selenium concentration of the culture medium with 10% FBS was 25.76 nM. The concentrations of selenium in the experiments are the supplemented concentrations, in addition to the selenium contained in the FBS; therefore, the culture medium of the control group (0 nM) contained 25.76 nM of selenium from the FBS.

SAMPLE PREPARATION FOR GPX-1 ACTIVITY ASSAY AND WESTERN BLOT ANALYSIS

After incubation in various concentrations of sodium selenite (0, 25, 50, 100, and 200 nM) for 72 h, cells were washed twice with phosphate-buffered saline and harvested; proteins were then extracted using RIPA buffer (Sigma, St. Louis, USA) with 10% protein inhibitor (Sigma, St. Louis, USA). Protein concentrations were determined using a Bio-Rad DC protein assay kit (Bio-Rad, Tokyo, Japan), following the method described by Lowry (Lowry *et al.*, 1951). The extracted sample was stored at -80°C until the GPx-1 activity assay and western blot analysis were performed.

DETERMINATION OF TOTAL SELENIUM CONCENTRATION

Before treatment with sodium selenite, concentrations of selenium in FBS were measured using a method described previously (Watkinson, 1966). Briefly, a 0.1-mL sample was digested in a heating block with 2 mL acid mixture (nitric acid/perchloric acid at 2:1). The temperature was gradually raised from 50°C to 190°C, and the sample was incubated overnight. Selenate was reduced to selenite using 0.5 mL of 10 N hydrochloric acid at 150°C for 20 min. The selenium concentration was calculated based on fluorometric measurements at an excitation wavelength of 378 nm and an emission wavelength of 525 nm to determine the concentration of piaselelol, which is produced by the reaction of selenite with 2,3-diaminonaphthalene. The accuracy of this analysis was monitored by the measurement of bovine liver SRM 1577b as the reference material (National

Institute of Standards and Technology, MD, USA).

MTT ASSAY

Selenium has a very narrow therapeutic dose window (Rayman, 2008); therefore, we confirmed the selenium dose and associated cell viability using a methyl thiazolyltetrazolium assay (MTT) assay, as described previously (Faried *et al.*, 2006). Cell proliferation inhibition analysis was performed in cells in the presence of various concentrations of sodium selenite and different incubation times. Briefly, cells (2×10^4 in 50 μ L/well) were plated in 96-well plates. After the initial cell seeding, sodium selenite was added in a concentration range of 0 to 10 μ M, and cells were incubated for 24, 48, and 72 h. Then, 10 μ L of WST-8 assay cell-counting solution (Dojindo Lab., Tokyo, Japan) was added to each well, and cells were then incubated at 37°C for 3 h. After the addition of 100 μ L/well of 1 N HCl, the cell proliferation rate was then determined by measuring the absorbance at a wavelength of 450 nm with a reference wavelength of 650 nm. The absorbance was read using a micro-titer plate reader (Becton-Dickinson, NJ, USA). The overall results were derived from four experiments.

GPx-1 ACTIVITY ASSAY

Based on a previous study (Hoefig *et al.*, 2011), we used 72 h as the optimal incubation time for sodium selenite supplementation. The enzymatic activity of GPx-1 was determined from Hc cell homogenates using the method described by Paglia and Valentine (1967). Briefly, the activity was indirectly monitored using spectrophotometric methods by observing the reduction of oxidized glutathione to reduced glutathione using nicotinamide adenine dinucleotide phosphate (NADPH) as the reducing agent. GPx-1 activity was quantified by measuring the change in NADPH absorbance at 340 nm and expressed as the change in NADPH (Δ mM NADPH) during the unit time (min) and mg protein in the presence of the substrate tert-Butyl hydroperoxide.

WESTERN BLOTS

Proteins (30 μ g and 40 μ g for ABCA-1) were electrophoresed on 5–20% polyacrylamide ready-made gels (Bio-Rad) and transferred onto a polyvinylidenedifluoride membrane (Millipore, Massachusetts, USA). HDL formation-related proteins were analyzed using the following antibodies: rabbit polyclonal anti-apoA-I (ab33470; Abcam, Cambridge, MA) at 1:200 dilution, goat polyclonal anti-apoA-II (178464; Calbiochem, Darmstadt, Germany) at 1:1000 dilution, rabbit polyclonal anti-GPx-1 (ab22604; Abcam) at 1:500 dilution, and rabbit polyclonal anti-ABCA-1 (NB400-105; Novus Biological, Littleton, USA) at 1:1000 dilution. Mouse monoclonal anti-GAPDH (MAB374; Abcam) antibody at 1:1000 dilution was used as the loading control. Secondary antibodies, namely donkey anti-rabbit IgG (NA934; Amersham, Buckinghamshire, UK), rabbit anti-goat IgG (ab7132; Abcam), or rabbit anti-mouse IgG (ab6728; Abcam), linked to horseradish peroxidase were then

applied, and immunoreactive bands were visualized using the prime chemiluminescence assay (ECL; Amersham). Scanning densitometry was performed by Image Quant LAS 4000 (Amersham), and autoradiographs were quantified using the National Institute of Health's ImageJ software program.

QUANTITATIVE REAL-TIME POLYMERASE CHAIN REACTION ANALYSIS

After incubation in various concentrations of sodium selenite (0, 25, 50, 100, and 200 nM) for 72 h, total RNA was isolated from Hc cells using TRIzol reagent (Invitrogen, CA, USA). RNA was prepared by reverse transcription using oligo-dT and dNTP, and each sample was processed with the RT-PCR kit (TAKARA, Japan). Quantitative real time-PCR was performed using the SYBR Green PCR Master Mix (Applied Biosystems, CA, USA) according to the manufacturer's instructions, and then evaluated using the LightCycler 480 Real-Time PCR system (Roche, CA, USA). The expression level of the target gene was normalized against GAPDH mRNA levels. The sequences of primers for quantitative real-time PCR used in this study are listed in Table 1.

Table 1- Primers for quantitative real-time PCR

	Forward	Reverse
hABC A-1	GAAGTGGCTGTGTT CCATGAT	GATGAGCCAGACT TCTGTTGC
hApo A-I	GCCTTGGGAAAAC AGCTAAACC	CCAGAAGTCTGG GTCACA
hApo A-II	CAAGAAGGCTGGA ACGGAAC	CTGGGGTTGGAA GACAATGG
hGAP DH	ACCACATCCATGCC ATCAC	TCCACCACCCTGT TGCTGTA
hGPx -1	CGCCAAGAACGAA GAGATTC	TCGATGTCAATGG TCTGGAA

hABCA-1, human ATP-binding cassette transporter-1; hApoA-I, human apolipoprotein A-I; hApoA-II, human apolipoprotein A-II; hGAPDH, human glyceraldehyde-3-phosphatase dehydrogenase; hGPx-1, glutathione peroxidase-1.

STATISTICAL ANALYSIS

Statistical analysis was performed using 1-way analysis of variance with Dunnett's post-hoc multiple comparison tests to compare between "no treatment" as the control group and each experimental group. A p-value < 0.05 was considered statistically significant. The statistical analysis of the data was performed with IBM SPSS (version 20.0 for Windows, IBM, NY, USA).

RESULTS

LOW-DOSE SODIUM SELENITE SUPPLEMENTATION DID NOT INHIBIT CELL PROLIFERATION

The Hc cells showed no inhibition of cell proliferation after 72-h incubation with low doses of sodium selenite; however, inhibition of cell proliferation

began to increase ($\geq 17.3\%$) at a concentration of 2.5 μM (Figure 1) and reached IC_{50} at 5 μM (Table 2).




Figure 1. Effects of sodium selenite supplementation on inhibition of cell proliferation
Data are presented as mean \pm SE and expressed as a percentage (%). The x-axis is presented in logarithmic scale. The results were derived from four experiments ($n = 8/\text{group}$).

Table 2. Summary of the time-resolved IC_{50} values of sodium selenite determined by MTT assay in Hc cells

Incubation times (h)	IC_{50} (μM)
24	-
48	7.0
72	5.0

Hc, human primary hepatocytes; IC_{50} , half-maximal inhibitory concentration; MTT assay, methyl thiazolyltetrazolium assay.

SODIUM SELENITE INCREASED GPx-1 ACTIVITY IN Hc CELLS

GPx-1 activity was measured in Hc cells after treatment with 0–200 nM sodium selenite (Figure 2). Sodium selenite supplementation increased GPx-1 activity significantly; this reached saturation levels at a concentration of 50 nM. Thus, GPx-1 activity was dependent on the presence of exogenous sodium selenite.




Figure 2- Effects of sodium selenite supplementation on GPx-1 activity
Data are presented as mean \pm SE, $n = 5/\text{group}$. **: $p < 0.01$ vs. control

EFFECTS OF SODIUM SELENITE SUPPLEMENTATION ON PRE β -HDL FORMATION-RELATED PROTEINS

Next, we observed the effects of sodium selenite supplementation on the expression of GPx-1 and

pre β -HDL formation-related proteins (apoA-I, apoA-II, and ABCA-1) in a dose range of 0 to 200 nM by western blotting and subsequent quantification by Image J (Figure 3). Sodium selenite supplementation significantly increased the protein expression of GPx-1 ($p < 0.01$ at 50 nM, and $p < 0.05$ at 100 nM) and apoA-I ($p < 0.05$ at 50 nM). These two protein expressions peaked at a dose of 50 nM. The expression of ABCA-1 also showed a tendency to increase although this was not significant. In contrast, selenium supplementation had no effects on apoA-II expression.




Figure 3. (a) Western blot analyses of GPx-1, apoA-I, apoA-II, and ABCA-1 of Hc cells after treatment with sodium selenite. (b) Quantification of western blot analyses using ImageJ software
(a) GAPDH expression was used as a reference. (b) Data are presented as mean \pm SE with $n = 6/\text{group}$. *: $p < 0.05$ vs. control. **: $p < 0.01$ vs. control.

EFFECTS OF SODIUM SELENITE SUPPLEMENTATION ON THE EXPRESSION OF PRE β -HDL FORMATION-RELATED TARGET GENES

The mRNA expressions of GPx-1 and several target genes related with pre β -HDL formation (apoA-I, apoA-II, and ABCA-1) were analyzed by real-time PCR (Figure 4). After supplementation with sodium selenite, a significant increase in the mRNA expression of GPx-1 ($p < 0.01$ at 50, 100, and 200 nM) was observed. Similarly, an upregulation in apoA-I mRNA expression was observed at 50 nM concentrations ($p < 0.05$). In contrast, no significant effect of sodium selenite supplementation was noted for ABCA-1 or apoA-II expression.




Figure 4. mRNA Expressions of GPx-1, apoA-I, apoA-II, and ABCA-1 in Hc cells after treatment with sodium selenite as analyzed by real-time PCR
Data are presented as mean \pm SE (n = 5/group). *: $p < 0.05$ vs. control. **: $p < 0.01$ vs. control.

DISCUSSION

In this study, we demonstrated that in Hc cells, GPx-1 activity and protein and mRNA expression were dependent on the presence of exogenous sodium selenite. Second, sodium selenite supplementation significantly increased protein and mRNA expressions of apoA-I, suggesting that selenium induced apoA-I expression from the transcription level. Interestingly, the peak of protein and mRNA expressions of GPx-1 and ApoA-I were reached at the same dose of sodium selenite (50 nM). Furthermore, ABCA-1 also showed a trend towards increased expression, although this was not significant. Third, selenium supplementation in the form of sodium selenite did not enhance apoA-II protein or mRNA expression.

In the present study, we used sodium selenite to induce GPx-1 activity and the expressions of related proteins and mRNA. In agreement with previous studies, we demonstrated that sodium selenite increased GPx-1 activity in Hc cells after 72-h incubation, with a dose saturation at 50 nM (Lei *et al.*, 1995; Hoefig *et al.*, 2011). Our results suggested the presence of a time lag between the addition of sodium selenite and the induction of GPx-1 synthesis. Furthermore, supplementation with selenium in the form of sodium selenite to normal hepatocyte cells increased the activity as well as the protein and mRNA expressions of GPx-1, thus protecting cells from potential damage by reactive oxygen species. Sodium selenite, an inorganic form of selenium, is well established as a dietary supplement, and its consumption has increased over the past few years (Rayman, 2008). In healthy people, all dietary requirements can be fulfilled solely from food sources; however, a recent study has reported that the enhancement of certain nutrients, including selenium, in the tissues is important (Monsen, 2000). In addition, antioxidant supplementation is

recommended for the prevention of oxidative damage in patients with chronic diseases (Actis-Goretta *et al.*, 2004).

Our study showed that sodium selenite supplementation in low doses ($<0.31 \mu\text{M}$) did not inhibit Hc cell proliferation after 72-h incubation. However, the proliferation of Hc cells was prominently inhibited at sodium selenite concentrations over $2.5 \mu\text{M}$. While selenium deficiency has been associated with several disorders, excessive selenium intake can cause selenium toxicity and even death (Rayman, 2008). It is therefore important to be aware of the possible implications of selenium supplementation in healthy people, especially for the prevention of CVD, as it can be beneficial, ineffectual, or even harmful for the human body, depending on the dosage.

In several clinical studies, supplementary selenium intake has been shown to have a positive correlation with HDL concentrations, leading to a beneficial effect in the prevention of CVD (Laclaustra *et al.*, 2010; Stranges *et al.*, 2011). Our study demonstrated that sodium selenite supplementation significantly increased the protein and mRNA expressions of apoA-I. Moreover, an upregulation trend in ABCA-1 protein expression was noted, although this was not statistically significant. These results suggest that selenium supplementation has an effect on pre β -HDL formation, inducing apoA-I expression from the transcription process.

Furthermore, selenium supplementation enhanced GPx-1 and apoA-I expressions in a concentration-dependent manner in Hc cells, with both expressions reaching their peaks at 50 nM. This dose was lower than the value reported in a previous study (Stahle *et al.*, 2009) using a human hepatoma cell line (hepG2), where the GPx-1 activity reached saturation at 100 nM of sodium selenite supplementation. Wilkening *et al.* (2003) concluded that hepatocytes are the most preferred *in vitro* models for assessing biotransformation in the human liver and for identifying compounds that are potentially toxic to humans. Considering that selenium has a very narrow therapeutic dose and that excessive selenium consumption can bring on toxicity and even death (Rayman, 2008), the exact concentrations of the supplemented dose of selenium and the type of cells used for examining the effects of selenium supplementation should be carefully considered.

Surprisingly, sodium selenite supplementation did not significantly affect apoA-II expression. This unexpected result suggests that this protein and gene may need stronger transcriptional inducers, such as oxidative stress. HDL concentrations depend on the following three major factors as follows: (1) production of apoA-I, apoA-II, and ABCA-1 by the liver; (2) the maturation of HDL; and (3) peripheral lipid docking to the liver (Panicker *et al.*, 2010). Our study merely focused on the first factor; therefore, the results do not incorporate all aspects of HDL production. Further *in vivo* studies are necessary to explore other potential roles of selenium in HDL formation.

CONCLUSION

In Hc cells that represent normal human liver cells, only a low dose of selenium is required to enhance the expressions of GPx-1 and apoA-I, a major component of pre β -HDL. In fact, higher doses of sodium selenite supplementation did not increase these upregulatory effects. Our findings suggest that selenium supplementation would be beneficial in preventing CVD events in healthy humans only when administered at the appropriatedose.

ACKNOWLEDGEMENTS

This study was supported by the GP 2012 grant (Gunma University, Japan). The authors would like to express their gratitude to Prof. Takeaki Nagamine for his kind gift of the Hc cells, and to ChihoYoshizawa and Miki Matsui for technical assistance.

REFERENCES

- Brenneisen P, Steinbrenner H, Sies H. Selenium, oxidative stress, and health aspects. *Mol Asp Med.* 2005; 26: 256–267.
- Abdulah R, Faried A, Kobayashi K, Yamazaki C, Suradji EW, Ito K, Suzuki K, Murakami M, Kuwano H, Koyama H. Selenium enrichment of broccoli sprout extract increases chemosensitivity and apoptosis of LNCaP prostate cancer cells. *BMC Cancer.* 2009; 9: 414.
- Brenneisen P, Steinbrenner H, Sies H. Selenium, oxidative stress, and health aspects. *Mol Asp Med.* 2005; 26: 256–267.
- Abdulah R, Faried A, Kobayashi K, Yamazaki C, Suradji EW, Ito K, Suzuki K, Murakami M, Kuwano H, Koyama H. Selenium enrichment of broccoli sprout extract increases chemosensitivity and apoptosis of LNCaP prostate cancer cells. *BMC Cancer.* 2009; 9: 414.
- Abdulah R, Miyazaki K, Nakazawa M, Koyama H. Chemical forms of selenium for cancer prevention. *J Trace Elem Med Biol.* 2005; 19: 141–150.
- Blankenberg S, Rupprecht HJ, Bickel C, Torzewski M, Hafner G, Tiret L, Smieja M, Cambien F, Meyer J, Lackner KJ, Investigators A. Glutathione peroxidase 1 activity and cardiovascular events in patients with coronary artery disease. *N Engl J Med.* 2003; 349: 1605–1613.
- WHO. Death and disability due to CVDs (heart attacks and strokes). In: Mendis S, Puska P, Norrving B (eds.). *Global atlas on cardiovascular disease prevention and control.* Geneva: World Health Organization in collaboration with the World Heart Federation and the World Stroke Organization. 2011; 2–13.
- Van der Steeg WA, Holme I, Boekholdt SM, Larsen ML, Lindahl C, Stroes ES, Tikkanen MJ, Wareham NJ, Faergeman O, Olsson AG, Pedersen TR, Khaw KT, Kastelein JJ. High-density lipoprotein cholesterol, high-density lipoprotein particle size, and apolipoprotein A-I: significance for cardiovascular risk: the IDEAL and EPIC-Norfolk studies. *J Am Coll Cardiol.* 2008; 51: 634–642.
- Ferrier H, Denise R. Plasma lipoproteins. In: Harvey RA (ed.). *Lippincott's illustrated reviews biochemistry 5th edition.* Baltimore: Lippincott Williams & Wilkins, a Wolter Kluwer business. 2011; 227–237.
- Kontush A, Chapman MJ. Normal functional high-density lipoprotein. In: *High-density lipoproteins structure, metabolism, function, and therapeutics.* New Jersey: John Wiley & Sons, Inc. 2012; 1–304.
- Miyazaki Y, Koyama H, Nojiri M, Suzuki S. Relationship of dietary intake of fish and non-fish selenium to serum lipids in Japanese rural coastal community. *J Trace Elem Med Biol.* 2002; 16: 83–90.
- Koyama H, Watanabe C, Satoh H, Hosokai H, Tamura S. Consistent relationship between selenium and apolipoprotein A-II concentrations in the sera of fasting middle-aged male abstainers and regular consumers of alcohol. *Biol Trace Elem Res.* 1995; 50: 33–42.
- Laclaustra M, Stranges S, Navas-Acien A, Ordovas JM, Guallar E. Serum selenium and serum lipids in US adults: National Health and Nutrition Examination Survey (NHANES) 2003-2004. *Atherosclerosis.* 2010; 210: 643–648.
- Stranges S, Tabák a G, Guallar E, Rayman MP, Akbaraly TN, Laclaustra M, Alftan G, Mussalo-Rauhamaa H, Viikari JS a, Raitakari OT, Kivimäki M. Selenium status and blood lipids: the cardiovascular risk in Young Finns study. *J Intern Med.* 2011; 270: 469–477.
- Stahle JA, Vunta H, Channa Reddy C, Sandeep Prabhu K. Regulation of expression of apolipoprotein A-I by selenium status in human liver hepatoblastoma cells. *Eur J Nutr.* 2009; 48: 283–290.
- Actis-Goretta L, Carrasquedo F, Fraga CG. The regular supplementation with an antioxidant mixture decreases oxidative stress in healthy humans. Gender effect. *Clin Chim Acta.* 2004; 349: 97–103.
- Colpo E, Vilanova CDDA, Brenner Reetz LG, Medeiros Frescura Duarte MM, Farias ILG, Irineu Muller E, Muller ALH, Moraes Flores EM, Wagner R, da Rocha JBT. A single consumption of high amounts of the Brazil nuts improves lipid profile of healthy volunteers. *J Nutr Metab.* 2013; 653185.
- Kaur HD, Bansal MP. Studies on HDL associated enzymes under experimental hypercholesterolemia:

possible modulation on selenium supplementation. *Lipids Health Dis.* 2009; 8: 55.

- Rees K, Hartley L, Day C, Flowers N, Clarke A, Stranges S. Selenium supplementation for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev.* 2013; 1: CD009671.
- Panicker S, Swathy SS, John F, Madambath I. Impact of Selenium on NF κ B Translocation in Isoproterenol-Induced Myocardial Infarction in Rats. *Biol Trace Elem Res.* 2010; 202–211.
- Wang X, Chen M, Li W, Zhu Y. ROS and NF-kappa B but not LXR mediate IL-1 beta signaling for the downregulation of ATP-binding cassette transporter A1. *Am J Physiol Cell Physiol.* 2006;1493–1501.
- Kretz-Remy C, Arrigo AP. Selenium: a key element that controls NF-kappa B activation and I kappa B alpha half life. *Biofactors.* 2001; 14: 117–125.
- Makropoulos V, Brüning T, Schulze-Osthoff K. Selenium-mediated inhibition of transcription factor NF-kappa B and HIV-1 LTR promoter activity. *Arch Toxicol.* 1996; 70: 277–283.
- Hayakawa K, Nagamine T. Fucoidan-dependent increased membrane components in HepG2 cells: effect of fucoidan is not due to gene expression. *Cancer Genomics Proteomics.* 2014;114:93–113.
- Lowry OH, Rosebrough NJ, Farr L, Randall RJ. Protein measurement with the folin phenol reagent. *J Biol Chem.* 1951; 193(1):265–275.
- Watkinson JH. Fluorometric determination of selenium in biological material with 2,3-diaminonaphthalene. *Anal Chem.* 1966; 38: 92–97.
- Rayman MP. Food-chain selenium and human health: emphasis on intake. *Br J Nutr.* 2008; 100: 254–268.
- Faried A, Faried LS, Kimura H, Sohda M, Nakajima M, Miyazaki T, Kato H, Kanuma T, Kuwano H. Differential sensitivity of paclitaxel-induced apoptosis in human esophageal squamous cell carcinoma cell lines. *Cancer Chemother Pharmacol.* 2006; 57: 301–308.
- Hoefig CS, Renko K, Köhrle J, Birringer M, Schomburg L. Comparison of different selenocompounds with respect to nutritional value vs. toxicity using liver cells in culture. *J Nutr Biochem.* 2011; 22: 945–955.
- Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med.* 1967; 70: 158–169.
- Lei XG, Evenson JK, Thompson KM, Sunde RA. Glutathione peroxidase and phospholipid hydroperoxide glutathione peroxidase are differentially regulated in rats by dietary selenium. *J Nutr.* 1995; 125: 1438–1446.
- Monsen ER. Dietary reference intakes for the antioxidant nutrients: vitamin C, vitamin E, selenium, and carotenoids. *J Am Diet Assoc.* 2000; 100: 637–640.
- Wilkening S, Stahl F, Bader A. Comparison of primary human hepatocytes and hepatoma cell line HepG2 with regard to their biotransformation properties. *Drug Metab Dispos.* 2003; 31: 1035–1042.